



Natural Medications for Psychiatric Disorders

David Mischoulon, MD, PhD

Director

Depression Clinical and Research Program

Massachusetts General Hospital

Joyce R. Tedlow Professor of Psychiatry

Harvard Medical School

Disclosures

My spouse/partner and I have the following relevant financial relationship with a commercial interest to disclose:

Organization	Support	Role
Nordic Naturals	Donated drug and placebo for clinical trial	Clinical investigator
heckel medizintechnik GmbH	Donated Whole Body Hyperthermia device for clinical trial	Clinical investigator
MGH Clinical Trials Network and Institute (CTNI)	Salary support through CTNI from multiple pharmaceutical companies and NIMH	Clinical Rater and Director of Education

Objectives



- To understand the evidence base for efficacy of natural therapies in psychiatry
- To identify the risks and benefits of various natural treatments in psychiatry
- To be able to educate patients in purchasing natural products in both over-the-counter and prescription forms

Pros and Cons of Natural Remedies

- In 2017, 30% of Americans used CAM practices and products (NCCIH/CDC, 2017)
 - About \$30.2 billion/year out-of-pocket cost
- Global market expected to grow from \$209 billion in 2017 to \$373 billion in 2025
- Easy access, good tolerability
- Used by many who don't respond to standard therapies
- Limited research, few rigorous studies
- “Natural” does NOT mean “safe”
- Toxicity, adverse effects, interactions
- Different preparations/purity
- Insurance does not cover them


Natural Antidepressants

- **St John's Wort:** ~40 clinical trials; monotherapy effective for depression; dosed at 900-1800 mg/day; watch out for photophobia and drug-drug interactions; don't combine with SSRIs (serotonin syndrome); little data on breastfeeding
- **SAME:** ~45 clinical trials; effective for depression as monotherapy and augmentation; dosed at 400-1600 mg/day (sometimes up to 3200 mg/day); no interactions; GI upset common; safe to combine with other medications; little data in pregnancy
- **Omega-3:** >30 trials in depression as monotherapy and augmentation; a few trials in bipolar disorder; best for depression or depressed phase of bipolar; dosed at 1-2 g/day (EPA:DHA = 3:2 preferred); safe combined with other meds; recent evidence suggests 4 g/day may be effective in inflammatory depression; probably safe in pregnancy




Natural Anxiolytic-Hypnotics

- **Kava:** >12 studies; effective for generalized anxiety; doses from 60-300 mg/day; recent negative 16-week study in N=171 with GAD, 120 mg bid (Sarris et al, 2020); cases of liver toxicity/death, but recent evidence suggests safety; use with caution and preferably for short periods
- **Valerian:** >35 studies; recent meta-analyses less supportive; effective for insomnia; doses from 450-600 mg at bedtime; few toxicity concerns; apparently safe in pregnancy and in elderly but caution advised
- **Melatonin:** ~20 studies, 2 strong meta-analyses; effective for insomnia, particularly if circadian disturbance-based; dosed at 0.3-5.0 mg/day; start low and increase gradually; some concerns about toxicity in immunosuppressed individuals; prolonged-release form (2mg) effective in elderly; effective in children



Second-Tier Antidepressants: 5-Hydroxy Tryptophan (5-HTP)




- Intermediate of L-tryptophan in production of serotonin
- Obtained from *Griffonia simplicifolia*
- Most studies >20 years old (interest in the serotonin hypothesis)
- After arrival of SSRIs (~1987), 5-HTP became less compelling
- Association with the Eosinophilia-Myalgia Syndrome (1989-1990)
- 1500 cases, at least 38 deaths; banned by FDA
- Contamination attributed to bacterial fermentation, poor filtration
- Current manufacturing methods unlikely to produce EMS

Efficacy and Safety

- About 27 published studies for depression
 - Samples small; only 6 showed statistical significance
 - Doses of 20-3250 mg/d, typically 200-300 mg/d
 - Dosing frequency 3-4X/day due to relatively short half-life (4.3 ± 2.8 hr)
 - Meta-analyses suggest only 1-2 rigorous studies
 - Most common adverse effects are gastrointestinal (nausea, vomiting, and diarrhea)
 - 5-HTP plus SSRI or MAOI may cause serotonin syndrome


Versatile Agents: Inositol

- Sugar alcohol, structural isomer of glucose, located primarily within cell membranes
- Present in beans, grains, nuts, and many fruits. Average adult consumes 1 g/day
- Also called Vitamin B8
- Vital in second messenger system for numerous neurotransmitter receptors



Mechanisms of Action

1. Involved in synthesis of membrane phospholipids
2. Precursor in phosphatidylinositol (PI) cycle



www.johnnysilva.com

Efficacy

- 6 clinical trials for depression (5 PBO-controlled)
 - 1 monotherapy, 5 augmentation
 - 2 MDD, 1 unipolar + bipolar depression, the rest bipolar depression
 - Inositol > placebo in 3 of 5 controlled studies
 - Small samples; significance reached in only one study
- Also effective for panic disorder, OCD, bulimia nervosa
- Possible broad spectrum of action similar to SSRIs
- Negative in schizophrenia, ADHD, Alzheimer's, autism, ECT-induced cognitive impairment

Safety and Tolerability

- Side effects: mild increases in plasma glucose, gas, nausea, sleep disturbance, dizziness, headache
- Case reports of mania in bipolar depression
- No reported toxicity or drug-drug interactions
- Not recommended for pregnant women, given risk of inducing uterine contractions
- Recommended doses between 6-20 g/d, usually 12 g/d divided 2-4X/day



Versatile Agents: Rhodiola Rosea

- Found in mountains of Europe and Asia
- Used for centuries in traditional medicine of Asia, Scandinavia, Eastern Europe
- “Adaptogen” -- increases resistance to chemical, biological, and physical stressors
 - Stimulates nervous system
 - Enhances physical and mental performance
 - Prevents altitude sickness
 - Alleviates fatigue, stress, depression, sexual dysf.

Efficacy and Mechanisms

- Studied in Russia and Scandinavia for >40 years
 - Most reports not yet translated to English
- Multiple active ingredients
 - Adaptogenics (rosavins, tyrosol), antioxidants (flavonoids), monoamine modulation, MAO-A and B inhibition, opioid-like effects
- ≥ 4 controlled trials support efficacy in depression and anxiety as well as cognition
 - Doses from 100-680 mg/day


Recent RCT of Rhodiola for MDD

- Mild-to moderate MDD; N=100; 12 weeks
 - Group A: Sertraline ('high dose') + PBO
 - Group B: Sertraline and Rhodiola 600 mg/day
 - Group C: Sertraline + Rhodiola 300 mg/day
- All groups had statistically significant reduction in HAM-D, BDI, and CGI
 - Improvement was significantly greater for group B versus groups C and A
- Higher doses of Rhodiola may be best

Gao et al. J Affect Disord 2020; 265:99-103.

Safety and Tolerability

- SFX mild, uncommon
 - Allergy, irritability, fatigue, unpleasant sensations, especially at high doses
 - Insomnia/vivid dreams; take early in day
 - Best on empty stomach, before meals
- No interactions with other drugs
 - Combined with TCAs; reduces TCA side effects
- No data on pregnancy or bipolar cycling
 - Use with caution



For Dementia: Ginkgo Biloba

- Cognition enhancer; slows cognitive decline
- Approx. 30 studies in Alzheimer's, mostly supportive
- Contains flavonoids and terpene lactones
- Stabilizes neuronal membranes, scavenges free radicals
- Meta-analyses and systematic reviews suggest efficacy (Weinmann et al, 2010; Brondino et al, 2013; Hashiguchi et al, 2015; Liu et al, 2019)
- Cholinesterase inhibitors somewhat more effective but not as well tolerated; may be combined (Mazza et al, 2006; Yancheva et al, 2009; Cornelli, 2010; Nasab et al, 2012; Canevelli et al, 2014)
- No clear preventive effects (Andrade et al, 2009)

Clinical Recommendations

- Suggested dose = 120-240 mg/day
- Best started early; full assessment may require 1 year
- No data on longer-term impact
- May alleviate antidepressant-induced sexual dysfunction
- Side effects: mild GI upset, headache, irritability, dizziness, seizures in epileptics
- Inhibits platelet activating factor (PAF); may cause bleeding in patients on anticoagulants or having surgery
 - No increased risk of bleeding, based on hemostatic outcomes in meta-analysis of 18 trials (Kellermann et al, 2011)
 - PAF inhibition may increase risk of bleeding in pregnancy; risk to breastfeeding infants unknown

For Dementia: Folic Acid

- Association between low folic acid and depression, dementia
- High dietary intake associated with lower risk of dementia
- Low folate associated with worse cognitive function, higher risk of cognitive decline
- Low folate associated with decreased chance of response to antidepressants
- May protect brain via decreased homocysteine



L-methylfolate/Deplin

L-methylfolate

vs.

Synthetic Folic Acid

Bypasses any polymorphisms




Willems et al. Br J Pharmacol. 2004;141:825-30.

Cerefolin

- Cerefolin
 - 5.6 mg L-methylfolate (metafolin)
 - 1 mg of vitamin B12 (cyanocobalamin)
 - 50 mg of vitamin B2 (riboflavin)
 - 5 mg of vitamin B6 (pyridoxine)
- Cerefolin NAC
 - With methylcobalamin 2mg, N-acetylcysteine 600mg (increases glutathione, reduces oxidative damage)
- Approved for treatment or prevention of vitamin deficiencies (need Rx)
- Used off-label for psychiatric indications, including depression and dementia

(McCadden and Hudson, 2010)

Vayacog (Lipicogen)



- Phosphatidylserine (PS)
 - Important in cell membrane function
 - Cognitive decline with decreased brain PS-DHA
- Lipicogen (DHA-enriched PS) 310mg
 - PS 100mg; DHA 19.5mg; EPA 6.5mg
- Vakhapova et al, 2010, 2014; N=157; 15 weeks
 - 300mg/d vs PBO; + 15wk follow-up at 100mg/d
 - PS-DHA associated with significant improvement in sustained attention and memory recognition; maintained in continuation

CBD: Applications

- Multiple sclerosis pain: Nabiximols (Sativex) oral aerosolized mist containing CBD + THC
 - Each spray delivers 2.7 mg THC + 2.5 mg CBD
 - Approved in Canada since 2005; also in Sweden
- Epilepsy: numerous clinical trials show CBD effective for certain childhood epilepsy disorders
 - Oral cannabidiol solution (Epidiolex) FDA-approved in June 2018 for Lennox-Gastaut syndrome and Dravet syndrome
 - 10-20 mg/Kg/day
- Limited data on other indications

Cannabis Products: Systematic Review

- Possible reduction of social anxiety
- Mixed (mainly positive) evidence for adjunctive use in schizophrenia
- Limited evidence in insomnia and PTSD
- No evidence of benefit for depression from high THC therapeutics or for CBD in mania
- Some potential efficacy for an oral cannabinoid/terpene combination in ADHD.

Sarris et al. BMC Psychiatry 2020; 20:24. doi: 10.1186/s12888-019-2409-8.

CBD: Mechanisms

- Indirect antagonist for CB1 and CB2 receptors
 - Potentiates THC by increasing CB1 receptor density or through other CB1 receptor-related mechanisms
- Interacts with G protein-coupled receptors
- Serotonin 5-HT_{1A} receptor partial agonist
 - antidepressant, anxiolytic effects?
- Allosteric modulator of μ - and δ -opioid receptors

CBD: Safety

- Common Side Effects
 - sleepiness
 - decreased appetite
 - diarrhea
 - fatigue
 - malaise
 - weakness
 - insomnia
- No intoxicating effects as with THC

CBD Recommendations

- Caution with high-THC formulations
 - esp. in youth, and anxiety or psychotic disorders
- Slow titration
- Regular assessment
- Caution in cardiovascular, respiratory disorders, pregnancy and breast-feeding
- Consider occupational safety as well

Sarris et al. BMC Psychiatry 2020; 20:24. doi: 10.1186/s12888-019-2409-8.

Light Therapy



- Best studied treatment for Seasonal Affective Disorder (SAD)
- Box containing fluorescent lamps
- 10,000 lux full spectrum or cool white fluorescent lights behind UV shield
- 30 min-2 hrs per day, for 2-4 weeks
- Lower-intensity (e.g. 2,500 lux) may require longer sessions

Efficacy

- Many studies, though most are small and not as rigorous as we would like
- Pooled analysis: 53% attain full remission
- 43% of cases of moderate-severe SAD attained remission
- BUT light therapy is often not enough
- May need to combine with antidepressants

Safety




- Side effects
 - Headache
 - Eyestrain
 - Agitation/Irritation
 - Nausea
- Managing side effects
 - Reduce treatment time
 - Move farther from light box
 - Take breaks between sessions
 - Change time of day of use
- Relative Contraindications
 - Retinal disease
 - Medications or medical conditions (e.g., Lupus) that increase light sensitivity
 - Mania or hypomania can be triggered by light therapy in cases of bipolar disorder

Other Indications for Light Therapy

- Types of depression/mood disorders that don't occur seasonally
- In chronic depression, add light to ongoing antidepressant treatment in fall/winter
- Jet lag (can buy portable lights for travel)
- Sleep disorders
- Adjusting to a nighttime work schedule
- Dementia



Proper Use



- Morning light may be more effective
- BID dosing may be best (AM and PM)
- Set up on desk or next to easy chair; use while reading, watching TV, working at computer, etc.
- Light should hit eyes *indirectly*
 - Direct light can harm eyes

When to Use the Lamp

- Begin in early fall, when it becomes cloudier
- Continue until spring, when outdoor light can sustain good mood and energy
- May also use off-season, e.g., during a cloudy spell
- But above all, find a schedule that works and stick with it



Exercise



- Aerobic exercise
 - Stationary bike, 2 sessions per day, 25 mins in afternoon, for one week
 - As effective as light Rx compared to controls
- Aerobic exercise under bright light (2500-4000 lux) is better than no exercise or exercise under normal light for atypical symptoms
- Outdoor exercise may therefore be best!

Synthesis:

CANMAT Recommendations

Table 3. Summary of Recommendations for Natural Health Products.

Intervention	Indication	Recommendation	Evidence	Monotherapy or Adjunctive Therapy
St. John's wort	Mild to moderate MDD	First line	Level 1	Monotherapy
	Moderate to severe MDD	Second line	Level 2	Adjunctive
Omega-3	Mild to moderate MDD	Second line	Level 1	Monotherapy or adjunctive
	Moderate to severe MDD	Second line	Level 2	Adjunctive
SAM-e	Mild to moderate MDD	Second line	Level 1	Adjunctive
	Moderate to severe MDD	Second line	Level 2	Adjunctive
Acetyl-L-carnitine	Mild to moderate MDD	Third line	Level 2	Monotherapy
<i>Crocus sativus</i> (saffron)	Mild to moderate MDD	Third line	Level 2	Monotherapy or adjunctive
DHEA	Mild to moderate MDD	Third line	Level 2	Monotherapy
Folate	Mild to moderate MDD	Third line	Level 2	Adjunctive
<i>Lavandula</i> (lavender)	Mild to moderate MDD	Third line	Level 3	Adjunctive
Inositol	Mild to moderate MDD	Not recommended	Level 2	
Tryptophan	Mild to moderate MDD	Not recommended	Level 2	
<i>Rhodiola rosea</i> (roseroot)	Mild to moderate MDD	Not recommended	Insufficient evidence	

DHEA, dehydroepiandrosterone; MDD, major depressive disorder; SAM-e, S-adenosyl-L-methionine.

Ravindran et al, Can J Psychiatry 2016; 61: 576-587

CANMAT Recommendations (cont'd)

Table 2. Summary of Recommendations for Physical and Meditative Treatments.

Intervention	Indication	Recommendation	Evidence	Monotherapy or Adjunctive Therapy
Exercise	Mild to moderate MDD	First line	Level 1	Monotherapy
	Moderate to severe MDD	Second line	Level 1	Adjunctive
Light therapy	Seasonal (winter) MDD	First line	Level 1	Monotherapy
	Mild to moderate nonseasonal MDD	Second line	Level 2	Monotherapy and adjunctive
Yoga	Mild to moderate MDD	Second line	Level 2	Adjunctive
Acupuncture	Mild to moderate MDD	Third line	Level 2	Adjunctive
Sleep deprivation	Moderate to severe MDD	Third line	Level 2	Adjunctive

MDD, major depressive disorder.

Ravindran et al, Can J Psychiatry 2016; 61: 576-587

Conclusions:

Who Should Use Natural Remedies?

- Mildly ill people with a strong interest in natural remedies who don't mind the cost
- People who have tried most everything else and have not responded, or had many side effects
 - But they are often the most difficult to treat
- Be careful with
 - Pregnant or breastfeeding women
 - Patients on multiple medications
 - Beware drug-drug interactions!

Questions?

Please email me at:

dmischoulon@mgh.harvard.edu